Lecture 2: General Equilibrium

Mauricio Romero

Lecture 2: General Equilibrium

Cobb-Douglas

Using calculus Perfect substitutes Perfect complements

$$u_A(x,y) = x^{\alpha} y^{1-\alpha}$$

$$u_B(x,y) = x^{\beta} y^{1-\beta}$$


For graph suppose

$$\alpha = 0.7$$

$$\beta = 0.3$$

$$\omega^{A} = (1, 1)$$

$$\omega^{B} = (1, 1)$$



- ▶ Indifference curves must be tangent (formalize this later)
- Thus, the MRS must be equalized across the two consumers

$$MRS_{x,y}^{A} = \frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}}$$

$$MRS_{x,y}^{B} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

$$\frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

- ▶ Indifference curves must be tangent (formalize this later)
- Thus, the MRS must be equalized across the two consumers

$$MRS_{x,y}^{A} = \frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}}$$

$$MRS_{x,y}^{B} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

$$\frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

But we haven't used the fact that

$$x^A + x^B = \omega_x$$
$$y^A + y^B = \omega_y$$

- Indifference curves must be tangent (formalize this later)
- ▶ Thus, the MRS must be equalized across the two consumers

$$MRS_{x,y}^{A} = \frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}}$$

$$MRS_{x,y}^{B} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

$$\frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

But we haven't used the fact that

$$x^A + x^B = \omega_x$$
$$y^A + y^B = \omega_y$$

$$\frac{\alpha}{1-\alpha} \frac{y^A}{x^A} = \frac{\beta}{1-\beta} \frac{\omega_y - y^A}{\omega_x - x^A}$$

- Indifference curves must be tangent (formalize this later)
- ► Thus, the MRS must be equalized across the two consumers

$$MRS_{x,y}^{A} = \frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}}$$

$$MRS_{x,y}^{B} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

$$\frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}} = \frac{\beta}{1 - \beta} \frac{y^{B}}{x^{B}}$$

But we haven't used the fact that


$$y^{A} + y^{B} = \omega_{y}$$

$$\frac{\alpha}{1 - \alpha} \frac{y^{A}}{x^{A}} = \frac{\beta}{1 - \beta} \frac{\omega_{y} - y^{A}}{\omega_{x} - x^{A}}$$

 $x^A + x^B = \omega_x$

$$y^{A} = \frac{1 - \alpha}{\alpha} \frac{\beta}{1 - \beta} \frac{\omega_{y} x^{A}}{\omega_{x} + \left(\frac{1 - \alpha}{\alpha} \frac{\beta}{1 - \beta} - 1\right) x^{A}}$$





Lecture 2: General Equilibrium

Cobb-Douglas
Using calculus
Perfect substitutes
Perfect complement

Using calculus

Essentially in this exercise we are doing the following:

$$\max_{(x^A,y^A),(x^B,y^B)}u_A(x^A,y^A) \text{ such that}$$

$$u_B(x^B,y^B)\geq \underline{u}_B=u_B(x^{B^*},y^{B^*})$$

$$x^B+x^A\leq \omega_x,$$

$$y^B+y^A\leq \omega_y.$$

Theorem

Consider an Edgeworth Box economy and suppose that all consumers have strictly monotone utility functions. Then a feasible allocation $(x^{A^*}, y^{A^*}, x^{B^*}, y^{B^*})$ is Pareto efficient if and only if it solves

$$\max_{(x^A,y^A),(x^B,y^B)} u_A(x^A,y^A)$$
 such that $u_B(x^B,y^B) \geq \underline{u}_B = u_B(x^{B^*},y^{B^*})$ $x^B + x^A \leq \omega_x,$ $y^B + y^A \leq \omega_y.$

Very tempting to use lagrangeans, no?

► We need to assume all consumers have quasi-concave, strictly monotone, differentiable utility functions

Then we can solve:

$$\max_{(x^A, y^A)} u_A(x^A, y^A) + \lambda (u_B(\omega_X - x^A, \omega_y - x^B) - \underline{u}_B).$$

Lets take the first order conditions of the above problem:

$$\frac{\partial u_{A}}{\partial x}(x^{A^{*}}, y^{A^{*}}) = \lambda \frac{\partial u_{B}}{\partial x}(\omega_{x} - x^{A^{*}}, \omega_{y} - x^{B^{*}})$$
$$\frac{\partial u_{A}}{\partial y}(x^{A^{*}}, y^{A^{*}}) = \lambda \frac{\partial u_{B}}{\partial y}(\omega_{x} - x^{A^{*}}, \omega_{y} - x^{B^{*}})$$

If $(x^{A^*}, y^{A^*}, x^{B^*}, y^{B^*})$ is Pareto efficient then

$$\frac{\frac{\partial u_A}{\partial x}(x^{A^*},y^{A^*})}{\frac{\partial u_A}{\partial y}(x^{A^*},y^{A^*})} = \frac{\frac{\partial u_B}{\partial x}(\omega_x - x^{A^*},\omega_y - y^{A^*})}{\frac{\partial u_B}{\partial y}(\omega_x - x^{A^*},\omega_y - y^{A^*})} = \frac{\frac{\partial u_B}{\partial x}(x^{B^*},y^{B^*})}{\frac{\partial u_B}{\partial y}(x^{B^*},y^{B^*})}.$$

▶ In short $MRS_{x,y}^A = MRS_{x,y}^B$

► This condition is *necessary* and *sufficient*

Theorem

Suppose that both consumers have utility functions that are quasi-concave and strictly increasing. Suppose that $(x^{A*}, y^{A*}, \omega_x - x^{A*}, \omega_y - y^{A*})$ is an **interior** feasible allocation. Then $(x^{A*}, y^{A*}, \omega_x - x^{A*}, \omega_y - y^{A*})$ is Pareto efficient if and only if

$$\frac{\frac{\partial u_A}{\partial x}(x^{A^*}, y^{A^*})}{\frac{\partial u_A}{\partial y}(x^{A^*}, y^{A^*})} = \frac{\frac{\partial u_B}{\partial x}(\omega_x - x^{A^*}, \omega_y - y^{A^*})}{\frac{\partial u_B}{\partial y}(\omega_x - x^{A^*}, \omega_y - y^{A^*})} = \frac{\frac{\partial u_B}{\partial x}(x^{B^*}, y^{B^*})}{\frac{\partial u_B}{\partial y}(x^{B^*}, y^{B^*})}.$$

Intuition

Suppose that we are at an allocation where $MRS_{x,y}^A=2>MRS_{x,y}^B=1$. Can we make both consumers better off?

Intuition

Suppose that we are at an allocation where $MRS_{x,y}^A=2>MRS_{x,y}^B=1$. Can we make both consumers better off?

- \blacktriangleright A gives up 1 unit of y to person B in exchange for unit of x
- ▶ *B* is indifferent since his $MRS_{x,y}^B = 1$.
- ➤ A receives a unit of x and only needs to give one unit of y (he was willing to give two)
- ► We have reallocated goods to make *A* strictly better off without hurting *B*

General case

$$\max_{((x_1^1,\ldots,x_L^l),\ldots,(x_1^l,\ldots,x_L^l))} u_1(x_1^1,\ldots,x_L^l) \text{ such that } u_2(x_1^2,\ldots,x_L^2) \geq \underline{u}_2,$$

$$\vdots$$

$$u_I(x_1^I,\ldots,x_L^I) \geq \underline{u}_I,$$

$$x_1^1+\cdots+x_I^I \leq \omega_1,$$

$$\vdots$$

$$x_I^1+\cdots+x_I^I \leq \omega_L.$$

General case

Theorem

Suppose that all utility functions are strictly increasing and quasi-concave. Suppose also that $((\hat{x}_1^1,\ldots,\hat{x}_L^1),\ldots,(\hat{x}_1^I,\ldots,\hat{x}_L^I))$ is a feasible interior allocation. Then $((\hat{x}_1^1,\ldots,\hat{x}_L^1),\ldots,(\hat{x}_1^I,\ldots,\hat{x}_L^I))$ is Pareto efficient if and only if $((\hat{x}_1^1,\ldots,\hat{x}_L^1),\ldots,(\hat{x}_1^I,\ldots,\hat{x}_L^I))$ exhausts all resources and for all pairs of goods ℓ,ℓ' ,

$$\mathit{MRS}^1_{\ell,\ell'}(\hat{x}^1_1,\ldots,\hat{x}^1_L) = \cdots = \mathit{MRS}^I_{\ell,\ell'}(\hat{x}^I_1,\ldots,\hat{x}^I_L).$$

▶ Utility functions must be strictly increasing, quasi-concave, and differentiable!

Lecture 2: General Equilibrium

Cobb-Douglas
Using calculus
Perfect substitutes
Perfect complement


Suppose that


$$u_A(x^A, y^A) = 2x^A + y^A$$


$$u_B(x^B, y^B) = x^B + y^B$$


$$\omega^A = (1, 1)$$


$$\omega^B = (1, 1)$$












Lecture 2: General Equilibrium


Cobb-Douglas
Using calculus
Perfect substitutes
Perfect complements


Perfect complements


Suppose that


$$u_A(x^A, y^A) = \min(x^A, 2y^A)$$
$$u_B(x^B, y^B) = \min(2x^B, y^B)$$
$$\omega^A = (3, 1)$$
$$\omega^B = (1, 3)$$
























































► What about: $u_A(x, y) = x^2 + y^2$, $u_B(x, y) = x + y$?

► Try it at home!

Recap

 We expect all exchanges to happen on the contract curve (hence its name)

▶ We expect all **voluntary** exchanges to be in the orange box

Can we say more? Not without prices